



Data on the use of dietary supplements in Danish patients with type 1 and type 2 diabetes

Ewers, B.; Trolle, Ellen; Jacobsen, S. S.; Vistisen, D.; Almdal, T. P.; Vilsbøll, T.; Bruun, J. M.

Published in:
Data in Brief

Link to article, DOI:
[10.1016/j.dib.2018.11.144](https://doi.org/10.1016/j.dib.2018.11.144)

Publication date:
2019

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Ewers, B., Trolle, E., Jacobsen, S. S., Vistisen, D., Almdal, T. P., Vilsbøll, T., & Bruun, J. M. (2019). Data on the use of dietary supplements in Danish patients with type 1 and type 2 diabetes. *Data in Brief*, 22, 241-244. <https://doi.org/10.1016/j.dib.2018.11.144>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Author's Accepted Manuscript

Data on the use of dietary supplements in Danish patients with type 1 and type 2 diabetes

B. Ewers, E. Trolle, S.S. Jacobsen, D. Vististen, T.P. Almdal, T. Vilsbøll, J.M. Bruun



PII: S2352-3409(18)31537-3S0899-9007(18)30231-4

DOI: <https://doi.org/10.1016/j.dib.2018.11.144>

Reference: DIB3559

To appear in: *Data in Brief*

Received date: 26 October 2018

Revised date: 29 November 2018

Accepted date: 30 November 2018

Cite this article as: B. Ewers, E. Trolle, S.S. Jacobsen, D. Vististen, T.P. Almdal, T. Vilsbøll and J.M. Bruun, Data on the use of dietary supplements in Danish patients with type 1 and type 2 diabetes, *Data in Brief*, <https://doi.org/10.1016/j.dib.2018.11.144>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Data Article

Title: Data on the use of dietary supplements in Danish patients with type 1 and type 2 diabetes

Authors: B. Ewers MSc¹, E. Trolle MSc², S.S. Jacobsen MSc¹, D. Vististen MSc, PhD¹, T.P. Almdal MD, DMSc³, T. Vilsbøll MD, DMSc^{1,4} and J.M. Bruun MD, PhD^{5,6}

Affiliations: ¹Steno Diabetes Center Copenhagen, Gentofte, Denmark ²The National Food Institute, Technical University of Denmark, Soeborg, Denmark, ³Rigshospitalet, Copenhagen, Denmark, ⁴Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark, ⁵Department of Nutrition, Exercise and Sports, Faculty of Science, University of Copenhagen, Frederiksberg, Denmark, ⁶Department of Medicine, Randers Regional Hospital, Denmark.

Contact email: *bettina.ewers@regionh.dk*

Abstract: The data in this article describes the use of dietary supplements in Danish patients with type 1 diabetes (T1D) and type 2 diabetes (T2D). The data was collected from a web-based dietary survey on dietary habits in 774 patients with T1D (n=426) and T2D (n=348). The data demonstrates that 99% of the patients with diabetes use dietary supplements with no gender differences. In comparison, only 64% in the general population use dietary supplements, "Dietary habits and adherence to dietary recommendations in patients with type 1 and type 2 diabetes compared with the general population in Denmark" [1].

A higher proportion of people in the general population use multivitamin/mineral supplementation as compared to patients with diabetes (48% vs. 34-37%) and a higher proportion of women than men with diabetes use multivitamin/mineral supplementation (T1D: 43% women vs. 26% men and T2D: 45% women vs. 34% men). More patients with diabetes than the general population use supplements such as calcium together with vitamin D, vitamin D, vitamin B, vitamin C, vitamin E, magnesium, calcium, Q10, ginger, garlic, and other herbal supplements.

Specifications Table

Subject area	<i>Nutrition</i>
More specific subject area	<i>Dietary supplementation in adults with type 1 and type 2 diabetes</i>
Type of data	<i>Table and text file</i>
How data was acquired	<i>Web-based questionnaire</i>
Data format	<i>Raw, analyzed</i>
Experimental factors	<i>The criteria used for including patients in the study and how data were collected has been described in Ewers et al. [1]</i>
Experimental features	<i>Data were analyzed by using SPSS software for Windows, version 22.0 (IBM Corp, Armonk, NY, USA)</i>
Data source location	<i>Copenhagen area, Denmark</i>
Data accessibility	<i>The data is with this article</i>
Related research article	Ewers B, Trolle E, Jacobsen SS, Vististen D, Almdal TP, Vilsbøll T, et al. Dietary habits and adherence to dietary recommendations in patients

	with type 1 and type 2 diabetes compared with the general population in Denmark. Nutrition (in press) [1].
--	--

Value of the data

- The data presented in this article provides new information about the use of dietary supplementation in patients with type 1 and type 2 diabetes
- The data provides important new evidence for differences in use of dietary supplementation between men and woman
- The data can be used to identify differences in the use of different dietary supplementation in patients with diabetes and the general population
- The data can be used by clinicians and academia for further research and as reference

Data

The article presents data on the intake of dietary supplementation in adult Danish patients with T1D and T2D. In **Table 1**, the percentage of dietary supplementation among male and female patients with T1D and T2D are presented. The supplementations are divided into all dietary supplements, multivitamin/mineral supplements, fish oil, calcium together with vitamin D, vitamin D and other supplements including herbal supplements. The data are compared with previous reported data on intake of dietary supplements in the general Danish population [2] based on the Danish National Survey of Dietary Habits in Denmark 2003-2008 [3].

Table 1. Percentage use of dietary supplementation among patients with diabetes and the general population

Dietary supplementation	T1D (n=426) %	T2D (n=348) %	General population (n=3,037) %
All dietary supplements,			
- Men	98.2	98.0	57.0
- Women	99.0*	100.0*	71.0
- All	98.6	98.6	64.0
Multivitamin/mineral,			
- Men	26.3	33.5	43.0
- Women	42.6*	44.6*	53.0
- All	34.0	37.0	48.0
Fish oil,			
- Men	20.3	21.8	18.0
- Women	27.3	24.8	25.0
- All	23.7	22.6	22.0
Calcium and vitamin D,			
- Men	11.1	10.9	5.2
- Women	30.1*	33.7*	17.9
- All	20.4	17.5	11.7
Vitamin D,			
- Men	19.8	27.0	6.4
- Women	25.4	43.6*	10.5
- All	22.5	31.8	8.5
Other supplements ¹ ,			

- Men	53.0	54.4	14.8
- Women	47.4*	37.6*	24.3
- All	50.2	49.6	19.8

¹Other supplements including vitamin B, C, E, magnesium, zinc, calcium, Q10, garlic, ginger, other herbal supplements.

*Statistical difference ($p < 0.05$) between men and women with diabetes.

Experimental Design, Materials and Methods

Data on use of dietary supplementation was collected in a cross-sectional dietary study of patients with T1D and T2D [1]. Data on the type of diabetes and gender was extracted from an electronic medical record in the outpatient clinic where the participants were recruited from. Participants were asked to report the use of all dietary supplements. Response to the use of the presented dietary supplements in Table 1 (yes/no) was mandatory. Furthermore, participants were asked to report the use of other dietary supplements (open-ended questions category). Data on the use of dietary supplements were compared between men and women with T1D and T2D by using the Chi-square test for differences in proportions. A two-sided significance level of $p < 0.05$ was used. All statistical analyses were performed with the SPSS software for Windows, version 22.0 (IBM Corp, Armonk, NY, USA).

Acknowledgements

The authors thank the patients with T1D and T2D in the outpatient clinic who participated in the survey and provided data for this article. We also thank the clinic at Steno Diabetes Center Copenhagen for funding.

References

- [1] Ewers B, Trolle E, Jacobsen SS, Vistisen D, Almdal TP, Vilsbøll T, et al. Dietary habits and adherence to dietary recommendations in patients with type 1 and type 2 diabetes compared with the general population in Denmark. *Nutrition* (in press)
- [2] Knudsen VK. Danskernes forbrug af kosttilskud. E-article. Technical University Denmark, Soeborg, Denmark, no 2, 2014
- [3] Pedersen AN, Fagt S, Groth MV, Christensen T, Biltoft-Jensen A, Matthiessen J, et al. Dietary habits in Denmark 2003-2008. Main results. The National Food Institute, Technical University Denmark. Soeborg, Denmark, 2010